



Toxicological Summary for: Thiamethoxam

CAS: **153719-23-4**

Synonyms: CGA 293343, 4H-1,3,5-Oxadiazin-4-imine, 3-[(2-chloro-5-thiazolyl)methyl] tetrahydro-5-methyl-N-nitro-, 3-((2-Chloro-5-thiazolyl)methyl)tetrahydro-5-methyl-N-nitro-4H-1,3,5-oxadiazin-4-imine

IUPAC name: 3-(2-chloro-1,3-thiazol-5-ylmethyl)-5-methyl-1,3,5-oxadiazinan-4-ylidene(nitro)amine

Acute Non-Cancer Health Based Value (nHBV_{Acute}) = Not Derived (Insufficient Data)

Short-term Non-Cancer Health Based Value (nHBV_{Short-term}) = 400 µg/L

(Reference Dose, mg/kg-d) x (Relative Source Contribution) x (Conversion Factor)
(Short-term Intake Rate, L/kg-d)

$$= \frac{(0.25 \text{ mg/kg-d}) \times (0.5)^* \times (1000 \text{ µg/mg})}{(0.285 \text{ L/kg-d})^{**}}$$

$$= 439 \text{ rounded to } \mathbf{400 \text{ µg/L}}$$

*Relative Source Contribution: MDH 2008, Section IV.E.1.

**Intake Rate: MDH 2008, Section IV.E.1. and US EPA 2011, Exposure Factors Handbook, Tables 3-1 and 3-81

Reference Dose/Concentration:	(POD x DAF)/Total UF = 0.25 mg/kg-d (Wistar rat)
Source of toxicity value:	Determined by MDH in 2016
Point of Departure (POD):	34.5 mg/kg-d (NOAEL, Brammer 2007)
Human Equivalent Dose (MDH, 2011):	POD x DAF = 34.5 mg/kg-d x 0.22 = 7.6 mg/kg-d
Total uncertainty factor (UF):	30
Uncertainty factor allocation:	3 for interspecies differences (for toxicodynamics), 10 for intraspecies variability
Critical effect(s):	Reduced pup body weight
Co-critical effect(s):	Hepatocyte hypertrophy, maternal death during pregnancy accompanied by hemorrhage of the uterus, bloody discharge in the perineal area, decreased number of animals with live fetuses, decreased fetal body weight, and increased fetal skeletal anomalies (fused sternebrae)
Additivity endpoint(s):	Developmental, Female Reproductive system, Hepatic (liver) system

Subchronic Non-Cancer Health Based Value (nHBV_{Subchronic}) = 200 µg/L

$$\frac{(\text{Reference Dose, mg/kg-d}) \times (\text{Relative Source Contribution}) \times (\text{Conversion Factor})}{(\text{Subchronic Intake Rate, L/kg-d})}$$
$$= \frac{(0.057 \text{ mg/kg-d}) \times (0.2)^* \times (1000 \text{ µg/mg})}{(0.070 \text{ L/kg-d})^{**}}$$
$$= 163 \text{ rounded to } \mathbf{200 \text{ µg/L}}$$

*Relative Source Contribution: MDH 2008, Section IV.E.1.

**Intake Rate: MDH 2008, Section IV.E.1. and US EPA 2011, Exposure Factors Handbook, Tables 3-1 and 3-81

Reference Dose/Concentration: (POD x DAF)/Total UF = 0.057 mg/kg-d (Beagles)
Source of toxicity value: Determined by MDH in 2016
Point of Departure (POD): 4.05 mg/kg-d (NOAEL, Altmann 1998)
Human Equivalent Dose (MDH, 2011): POD x DAF = 4.05 mg/kg-d x 0.43 = 1.7 mg/kg-d
Total uncertainty factor (UF): 30
Uncertainty factor allocation: 3 for interspecies differences (for toxicodynamics), 10 for intraspecies variability
Critical effect(s): Seminiferous tubule atrophy
Co-critical effect(s): None
Additivity endpoint(s): Male Reproductive system

Chronic Non-Cancer Health Based Value (nHBV_{Chronic}) = nHBV_{Subchronic} = 200 µg/L

$$\frac{(\text{Reference Dose, mg/kg-d}) \times (\text{Relative Source Contribution}) \times (\text{Conversion Factor})}{(\text{Chronic Intake Rate, L/kg-d})}$$
$$= \frac{(0.057 \text{ mg/kg-d}^{***}) \times (0.2)^* \times (1000 \text{ µg/mg})}{(0.044 \text{ L/kg-d})^{**}}$$
$$= 259 \text{ rounded to } \mathbf{300 \text{ µg/L}}$$

*Relative Source Contribution: MDH 2008, Section IV.E.1.

**Intake Rate: MDH 2008, Section IV.E.1. and US EPA 2011, Exposure Factors Handbook, Tables 3-1 and 3-81

***The calculated chronic RfD (0.43 mg/kg-d) is higher than the subchronic RfD (0.057 mg/kg-d), which is based on male reproductive effects. The chronic RfD must be protective of all types of adverse effects that could occur as a result of chronic exposure, including subchronic effects (MDH 2008). Therefore, the chronic RfD is set to the subchronic RfD. See the subchronic information above for details about the reference dose.

The Chronic nHBV must be protective of the acute, short-term, and subchronic exposures that occur within the chronic duration; and therefore, the Chronic nHBV is set equal to the Subchronic nHBV of 200 µg/L. Additivity endpoints: Male Reproductive system

Cancer Health Based Value (cHBV) = Not Derived

Cancer classification: Not likely to be carcinogenic to humans
Slope factor (SF): Not Applicable
Source of cancer SF: Not Applicable
Tumor site(s): Not Applicable

Volatile: No

Summary of Guidance Value History:

A pesticide rapid assessment of 20 µg/L was completed for thiamethoxam in 2014 by MDH. The 2016 nHBVs of 200 µg/L are higher than the pesticide rapid assessment due to the conservative method used for rapid assessments (MDH 2014). MDH intends to re-evaluate guidance values on a five year cycle in order to keep guidance values current with scientific knowledge. Under this process, thiamethoxam will undergo re-evaluation in 2021.

Summary of toxicity testing for health effects identified in the Health Standards Statute (144.0751):

Even if testing for a specific health effect was not conducted for this chemical, information about that effect might be available from studies conducted for other purposes. MDH has considered the following information in developing health protective guidance.

	Endocrine	Immunotoxicity	Development	Reproductive	Neurotoxicity
Tested for specific effect?	No	No	Yes	Yes	Yes
Effects observed?	Yes ¹	Yes ²	Yes ³	Yes ⁴	Yes ⁵

¹ A comprehensive toxicity study, specific for endocrine effects after thiamethoxam exposure, has not been completed; however, endocrine effects were observed in other studies. Endocrine effects included changes to the adrenal cortex and thyroid. In short-term and subchronic durations, adrenal gland changes occurred in male and female rats at thiamethoxam levels 400 to 700 times higher than the corresponding duration's reference dose. Thyroid gland changes occurred in rats between 200 - 700 times higher than the short-term reference dose.

² Immunological effects observed in thiamethoxam studies include changes in the thymus, spleen, and white blood cells. Changes to the thymus in rats were varied, with a range of changes between 250 to 1,400 times higher than the reference dose in short-term and subchronic durations. Conversely, one report observed no changes to the thymus at levels up to 690 times higher than the short-term reference dose. Changes to the spleen, in rat, were observed between levels of thiamethoxam 700 to 1,400 times higher than the subchronic reference dose. Beagles were a more sensitive species, with thymus and spleen changes occurring at levels 72 times higher than the short-term reference dose. White blood cell changes occurred at levels around 400 times higher than the subchronic reference dose.

³ The short-term reference dose is based on the developmental effect of reduced pup body weight, an effect that occurred in multiple studies. Reductions in fetal body weights and skeletal abnormalities

occurred at levels 700 times higher than the short-term reference dose in rats and 300 times higher than the short-term reference dose in rabbits.

⁴ The subchronic reference dose is based on adverse reproductive effects occurring in adult male beagles. There were no reproductive effects in pregnant rats exposed to levels of thiamethoxam up to 700 times higher than the short-term reference dose. Pregnant rabbits were more sensitive, demonstrating reproductive effects at levels 300 times higher than the short-term reference dose. Adult female mice experienced reproductive changes at levels up to 1,400 times higher than the subchronic reference dose, and male mice at levels over 3,000 times higher than the subchronic reference dose. Studies in adult rats varied, with reports of no adverse reproductive events at thiamethoxam levels 500 times higher than the subchronic reference dose, to changes in sperm concentrations at levels 16 times higher than the subchronic reference dose.

⁵ Thiamethoxam exposure in rodents produced transient neurotoxicity at high doses over 250 - 500 times greater than the short-term and subchronic reference doses. Altered activity and changes in the brain were noted in these studies. The mouse was more sensitive than the rat, with adverse effects in the mouse (reduced locomotor activity, convulsions, prostration) occurring at thiamethoxam levels 250 times higher than the short-term reference dose. The rat experienced adverse effects from thiamethoxam levels beginning at levels 450 times higher than the short-term reference dose. There were no adverse effects on pups when pregnant rats were exposed to thiamethoxam at levels up to 300 times higher than the short-term reference dose.

Resources Consulted During Review:

Altmann, B. (1996a). CGA 293343 technical - 3 month subchronic dietary toxicity study in Beagle dogs. Report No. 942107. Ciba-Geigy Ltd, Short/Long-term Toxicology, Stein, Switzerland. Unpublished, submitted to WHO by Syngenta Crop Protection AG.

Altmann, B. (1996b). CGA 293343 technical - 28 day range finding study in Beagle dogs. Study No. 942106. Novartis Crop Protection AG, Short/Long-term Toxicology, Basel, Switzerland. Unpublished.

Altmann, B. (1998). CGA 293343 technical - 12 month chronic dietary toxicity study in Beagle dogs. Report No. 942108. Novartis Crop Protection AG, Toxicology, Stein, Switzerland. Unpublished, submitted to WHO by Syngenta Crop Protection AG.

Australian Department of Health. Acceptable Daily Intakes for Agricultural and Veterinary Chemicals. Retrieved from www.health.gov.au

Bachmann, M. (1994). CGA 293343 technical - 28 Day exploratory toxicity study in male rats (gavage). Study No. 932107. Novartis Crop Protection, Inc., Short/long-term Toxicology, Stein, Switzerland. Unpublished.

Bachmann, M. (1995). CGA 293343 technical - 28 day range finding study in rats (administration in food). Report No. 942088. Ciba-Geigy Ltd, Short/Long-term Toxicology, Stein, Switzerland. Unpublished, submitted to WHO by Syngenta Crop Protection AG.

Bachmann, M. (1996a). CGA 293343 technical - 3 month oral toxicity study in rats (administration in food). Report No. 942089. Ciba-Geigy Ltd, Short/Long-term Toxicology, Stein, Switzerland. Unpublished, submitted to WHO by Syngenta Crop Protection AG.

- Bachmann, M. (1996b). CGA 293343 technical - 3 month range finding toxicity study in mice (administration in food). *Report No. 942105*. Ciba-Geigy Ltd, Short/Long-term Toxicology, Stein, Switzerland. Unpublished, submitted to WHO by Syngenta Crop Protection AG.
- Bachmann, M. (1998a). CGA 293343 technical - 18 month oncogenicity study in mice. Report No. 942109. Novartis Crop Protection AG, Toxicology, Stein, Switzerland. Unpublished, submitted to WHO by Syngenta Crop Protection AG.
- Bachmann, M. (1998b). CGA 293343 technical - 24 month carcinogenicity and chronic toxicity study in rats. Report No. 942110. Novartis Crop Protection AG, Experimental Toxicology, Stein, Switzerland. Unpublished, submitted to WHO by Syngenta Crop Protection AG.
- Brammer, A. (2003). Thiamethoxam: Preliminary Developmental Neurotoxicity Study in Rats. Report No. RR0935. Central Toxicity Laboratory, Alderley Park, Macclesfield, Cheshire, UK. Unpublished.
- Brammer, A. (2007). Thiamethoxam: Supplement to developmental neurotoxicity study in rats. Laboratory report No. RR0936-REG-S1. Central Toxicology Laboratory, Alderley Park, Macclesfield, Cheshire, UK. Unpublished, submitted to WHO by Syngenta Crop Protection AG.
- Brammer A. (2003). Thiamethoxam: Developmental neurotoxicity study in rats. Report No. RR0936. Central Toxicology Laboratory, Alderley Park, Macclesfield, Cheshire, UK. Unpublished, submitted to WHO by Syngenta Crop Protection AG. Retrieved from <http://www.epa.gov/pesticides/chemicalsearch/chemical/foia/cleared-reviews/reviews/060109/060109-2005-10-24a.pdf>
- California EPA. (2008). Summary of Toxicology Data: Thiamethoxam. Retrieved from <http://www.cdpr.ca.gov/docs/risk/toxsums/pdfs/5598.pdf>
- Dobrovetzky, M. (1998). CGA 293343 technical - rat dietary two-generation reproduction study, includes: Effects on sperm cell parameters. Report No. 942121 and 982015. Novartis Crop Protection AG, Toxicology, Stein, Switzerland. Unpublished, submitted to WHO by Syngenta Crop Protection AG.
- European Commission. (2007). European Union Pesticides Database. Retrieved from <http://ec.europa.eu/food/plant/pesticides/eu-pesticides-database/public/?event=activesubstance.detail&language=EN&selectedID=1937>
- European Commission. (2012). Directive 98/8/EC concerning the placing biocidal products on the market, Inclusion of active substances in Annex I or IA to Directive 98/8/EC, Assessment Report: Thiamethoxam Product-type 18. Retrieved from <https://circabc.europa.eu/sd/a/94c7a2fe-e407-411e-94cb-eadbb3823cd1/CA-Sept12-Doc%203%208a%20-%20AR%20Thiamethoxam%20PT18%20post%20SCB.pdf>
- McGregor, D., & Solecki, R. (2010). Pesticide residues in food - 2010. Joint FAO/WHO Meeting on Pesticide Residues, 565-676. Retrieved from http://apps.who.int/iris/bitstream/10665/44660/1/9789241665261_eng.pdf
- Minnema, D. (1997). Acute neurotoxicity study of orally administered CGA 293343 tech. in rats. Report No. 6117-364. Covance Laboratories Inc., Vienna, VA, USA. Unpublished, submitted to WHO by Syngenta Crop Protection AG.

- Minnema Daniel J. (1998). 13-Week Dietary Subchronic Neurotoxicity Study with CGA-293343 TECH in Rats. Unpublished by Covance Laboratories, Inc. Retrieved from <http://www.epa.gov/pesticides/chemicalsearch/chemical/foia/cleared-reviews/reviews/060109/060109-1999-06-25a.pdf>
- Minnesota Department of Health (MDH). (2008). Statement of Need and Reasonableness (SONAR), July 11, 2008. Support document relating to Health Risk Limits for Groundwater Rules. Retrieved from <http://www.health.state.mn.us/divs/eh/risk/rules/water/hrlsonar08.pdf>
- Minnesota Department of Health (MDH). (2011). MDH Health Risk Assessment Methods to Incorporate Human Equivalent Dose Calculations into Derivation of Oral Reference Doses. Retrieved from <http://www.health.state.mn.us/divs/eh/risk/guidance/hedrefguide.pdf>
- Minnesota Department of Health (MDH). (2014). Report on Pesticide Rapid Assessments. Retrieved from: <http://www.health.state.mn.us/divs/eh/risk/guidance/dwec/rapassrept.pdf>
- Moxon, M. (2004). Thiamethoxam: Two generation reproduction study in rats. Report No. CTL/RR0941/regulatory/report. Syngenta Ltd, Central Toxicology Laboratory, Alderley Park, Macclesfield, Cheshire, England. Unpublished, submitted to WHO by Syngenta Crop Protection AG.
- Oda, S. (1996a). An acute oral toxicity of CGA 293343 technical in rats. Report No. B-3120, CG 942111. Bozo Research Center Inc., Tokyo, Japan. Unpublished, submitted to WHO by Syngenta Crop Protection AG.
- Oda, S. (1996b). An acute oral toxicity study of CGA 293343 technical in mice. Report No. B-3122. Bozo Research Center Inc., Tokyo, Japan. Unpublished, submitted to WHO by Syngenta Crop Protection AG.
- Pastoor, T., Rose, P., Lloyd, S., Peffer, R., & Green, T. (2005). Case study: weight of evidence evaluation of the human health relevance of thiamethoxam-related mouse liver tumors. *Toxicol Sci*, 86(1), 56-60. doi:10.1093/toxsci/kfi126
- Suarez, F. A. (2009). Thiamethoxam. Syngenta Response to the Proposed Inclusion of Thiamethoxam on the List of Chemicals Submitted to the Carcinogen Identification Committee (March 2009). Syngenta Crop Protection, Inc. Retrieved from <http://oehha.ca.gov/media/downloads/proposition-65/crn/comments/syngentathiamethoxam.pdf>
- Swenberg, J. A. (1993). Alpha 2u-globulin nephropathy: review of the cellular and molecular mechanisms involved and their implications for human risk assessment. *Environ Health Perspect*, 101 Suppl 6, 39-44.
- U.S. Environmental Protection Agency - Office of Research and Development. (1988). Recommendations for and Documentation of Biological Values for Use in Risk Assessment. Retrieved from <http://cfpub.epa.gov/ncea/cfm/recordisplay.cfm?deid=34855>
- U.S. Environmental Protection Agency - Office of the Science Advisor. (2011). Recommended Use of Body Weight^{3/4} as the Default Method in Derivation of the Oral Reference Dose. Retrieved from <http://www.epa.gov/raf/publications/pdfs/recommended-use-of-bw34.pdf>

- U.S. Environmental Protection Agency - Office of Research and Development. (2011). Exposure Factors Handbook: 2011 Edition. Retrieved from <https://cfpub.epa.gov/ncea/risk/recordisplay.cfm?deid=236252>
- U.S. Environmental Protection Agency. (2010). Thiamethoxam - Human Health Risk Assessment for New Seed Treatment Uses on Alfalfa and Peanuts, and Use in Food Handling Establishments. Retrieved from <http://www.regulations.gov/#!documentDetail;D=EPA-HQ-OPP-2010-0602-0008>
- U.S. Environmental Protection Agency. (2011). Thiamethoxam. Human Health Assessment Scoping Document in Support of Registration Review. Retrieved from <http://www.regulations.gov/#!documentDetail;D=EPA-HQ-OPP-2011-0581-0005>
- U.S. Environmental Protection Agency. (2015). Human Health Benchmarks for Pesticides. Retrieved from <http://iaspub.epa.gov/apex/pesticides/f?p=HHBP:home>
- Winkler, G. (1996a). CGA 293343 technical - rabbit oral teratogenicity. Report No. 942119. Ciba-Geigy Ltd, Short/Long-term Toxicology, Stein, Switzerland. Unpublished, submitted to WHO by Syngenta Crop Protection AG.
- Winkler, G. (1996b). CGA 293343 technical - rat oral teratogenicity study. Report No. 942118. Ciba-Geigy Ltd, Short/Long-term Toxicology, Stein, Switzerland. Unpublished, submitted to WHO by Syngenta Crop Protection AG.