



*Protecting, maintaining and improving the health of all Minnesotans*

# **MINNESOTA EARLY HEARING DETECTION AND INTERVENTION (EHDI) PROGRAM**

## **Guideline for Organization and Administration of Universal Newborn Hearing Screening Programs in the Well Baby Nursery**

**Approved May, 2008**

### **Introduction**

The goal of an Early Hearing Detection and Intervention (EHDI) Program is to identify and provide effective intervention for newborns and infants with permanent childhood hearing loss and those at risk for hearing loss that may affect health, communication, learning and development. To achieve this goal, Minnesota law (Minnesota Statutes 144.966) requires that a hearing screen be performed on all newborns prior to hospital discharge or within one month if not a hospital birth. National standards specify that screening should be completed as soon as possible, and no later than one month of age; hearing loss should be clinically diagnosed as soon as possible, and no later than 3 months of age and intervention initiated as soon as possible, and no later than 6 months of age. With prompt referral and follow-up, Minnesota children have an opportunity to receive appropriate care and services even earlier.

Because of the importance of early identification of hearing loss, all screening, follow-up, and tracking procedures must at a minimum be consistent with current Minnesota statutory requirements. This document provides recommended guidelines for newborn hearing screening programs for infants in the Well Baby Nursery. Please see separate document regarding hearing screening guidelines for infants in the Special Care Nursery and Neonatal Intensive Care Unit (NICU). Additional resources are available from the Minnesota Department of Health (MDH) to assist hospitals and hearing screeners with specific issues of program development and management such as training, supervision, equipment options, and quality assurance issues ([www.health.state.mn.us/newbornscreening](http://www.health.state.mn.us/newbornscreening)).

### **Child and Family-Centered Communication**

Minnesota statute requires hospitals to present information to parents that covers the following topics:

- Potential risks and effects of hearing loss
- Benefits of early detection and intervention
- Nature of the screening procedure
- Applicable costs of screening procedure
- Parental option to have screening performed but to request that test results be destroyed by the MDH
- Parental option to refuse screening

Best practice would also include providing the following to parents:

- Prevalence of and risk factors for permanent childhood hearing loss (Appendix A)
- Possibility of late or progressive onset of hearing loss, including otitis media
- Developmental milestones for speech, language, and hearing (Appendix B)

### **Personnel**

Screening may be performed by trained personnel including the following:

- Audiologists, audiological technicians/assistants
- Nurses
- Nursing assistants
- Other trained medical personnel

Although licensed audiologists do not need to conduct the actual hearing screening, audiologists are uniquely qualified to develop and implement all aspects of an EHDI program. Hospital screening programs benefit from direct access to audiological consultation to address screening criteria, quality assurance, follow-up assessment and intervention services.

### **Training**

Training qualified screeners is an ongoing process. Training typically includes three phases: initial training and demonstration of skills, ongoing quality assurance, and refresher training. Resources for training may include experienced screening program managers and/or local licensed clinical and educational audiologists.

Initial training content is based on current best practice procedures as reported in professional literature. Assuring competent screeners is a critical component for every screening program. Training components include the following:

- Competency-based demonstration of trainees' performance in the nursery environment;

- Periodic observation of each screener in the nursery environment by a skilled professional such as an audiologist and/or program manager;
- Annual staff refresher training, with individual training available as needed.

## **Equipment**

Screening programs must use objective physiological screening methods such as Automated Auditory Brainstem Response (AABR) or otoacoustic emissions (DPOAEs or TOAEs) that do not require a behavioral response from the infant. OAE and AABR have proven to be effective screening measures. All equipment must meet technical specifications, calibration standards, and hospital safety standards.

## **Stimulus Parameters**

TOAEs should be measured in response to a click at approximately 80dB SPL (78-82dB SPL). DPOAEs should be measured in response to a series of paired tones (f1 and f2), with a ratio of 1.22 at a moderate level, where L1/L2= 65/55dB SPL.

Newborn screening AABRs typically are evoked using click stimuli at 30 to 35dB nHL at a moderate stimulus rate. Non-automated ABR testing is NOT recommended for newborn hearing screening programs in hospital nurseries due to issues of potential operator error and significant time/cost effectiveness issues.

Default stimulus parameters of either OAE or AABR equipment should be reviewed by skilled professionals such as a consulting audiologist or MDH audiologist to assure they are appropriate or to adjust them to be in accordance with clinically accepted national practices.

## **Pass/Refer Criteria**

### ***OAE***

Manufacturer's pass/fail criteria should be reviewed by the consulting audiologist or MDH audiologist and should be in accordance with clinically accepted national practices. Typical passing criteria for TOAE's include overall reproducibility greater than 50%, at least 50 low noise samples collected, stimulus stability of 75% or greater, and responses present at least 6dB above noise floor for at least three of the five test frequencies, with 4000Hz a mandatory passing frequency. Typical passing criteria for DPOAEs requires absolute response amplitude of at least -6dB and responses at least 6dB above the noise floor at three or more of the test frequency bands, with the 4000Hz region a mandatory passing frequency.

### ***AABR***

- Screening AABR pass criteria for newborns typically require repeatable Wave V evoked responses to clicks at  $\leq 35$  dB nHL for each ear, within specific latency

parameters. Once again the consulting audiologist or MDH audiologist should review the default pass/fail criteria for the specific AABR equipment used to ensure compliance with accepted national practices.

A quality screening program benefits from incorporating new and improved evidence-based technologies and procedures as they become available.

## **Hearing Screening Protocol**

The purpose of a screening test is to identify those infants at risk for hearing loss who need further testing. A screening test is not a diagnosis. A referral rate of 4% or less in the hospital nursery is considered acceptable.

### **INITIAL HEARING SCREENING**

The initial hearing screening is the first hearing screening performed on an infant after birth. The initial screening is preferably performed 12 hours or more after birth. The screening may be performed sooner if needed; however, a higher referral rate may occur due to residual birthing debris in the ear canal. For infants who fail the initial screen, hospitals should attempt to re-screen the infant prior to discharge. Inpatient hearing screening will consist of no more than two attempts using the same screening technique on each ear, assuming the infant is in an appropriate state for testing and there are neither equipment problems nor environmental interference during the test. The likelihood of obtaining a pass by chance alone is increased when screening is performed repeatedly.

If an infant does not pass the hearing screening in one or both ears, hospital and medical staff are strongly encouraged to immediately schedule a follow-up hearing appointment.

### **Documentation for Initial Hearing Screen**

Minnesota Statute 144.966 requires the following documentation:

- Screening results need to be recorded in the infant's medical record.
- Screening results need to be communicated to the parents of the infant so that they are well-understood.
- Screening results must be communicated to the infant's primary care physician in writing.
- Screening results must be reported to the MDH.

In order to improve follow-up for infants with failed hearing screens:

- Families of infants who refer on the hearing screening should be provided with information about why their baby may not have passed the hearing screening and

the importance of follow-up. In addition, a follow-up appointment for hearing re-screening or pediatric audiological evaluation should be scheduled.

- For infants who refer, the primary care physician and MDH should be promptly notified of the follow-up hearing appointment to facilitate timely services.

## **OUTPATIENT RE-SCREENING**

If the infant does not pass the hospital hearing screening in one or both ears, she or he needs to have a re-screen completed as soon as possible and no later than one month of age. It is ideal to allow 1-2 weeks for resolution of transient ear conditions to resolve before re-screening. Best practice would suggest that the following be included as part of the re-screening.

- A complete history should be taken, including neonatal history, family history of childhood hearing loss, and the parents' observations of their baby's response to sounds at home.
- Either AABR or OAE can be used for re-screening, regardless of which technology was used for the hospital hearing screening.
- Both ears are re-screened even if only one ear failed the initial screening.
- The re-screen should consist of a maximum of two attempts on each ear, assuming the infant is appropriately quiet for valid testing and there are neither equipment problems nor environmental interference during the test.

## **Documentation for Outpatient Re-Screening**

- Screening results must be recorded in the infant's medical record.
- Screening results need to be clearly communicated to the parents of the infant.
- Screening results must be communicated to the infant's primary care physician in writing.
- Screening results are reported to the MDH as required by MN Statute 144.966.
- Families of infants who refer on the hearing re-screening will be provided with information about why their baby may not have passed the hearing screening and the importance of follow-up. In addition, a diagnostic audiology appointment should be scheduled.
- For infants who refer, the primary care physician and the MDH should be promptly notified of the diagnostic audiology appointment to support timely services.

## **TRANSFERS**

If an infant is transferred, the hospital that discharges the infant home is responsible for screening

## **NON-HOSPITAL BIRTHS**

According to MN statute 144.966, a physician, nurse, midwife, or other health professional attending a birth outside a hospital or institution shall provide information, orally and in writing, as established by the Department of Health, to parents regarding places where the parents may have their infant's hearing screened and the importance of the screening.

## **RE-ADMITS**

Infants re-admitted to the hospital during the first month of life with conditions associated with potential hearing loss (e.g., hyperbilirubinemia, meningitis, sepsis) need to have the hearing screen repeated prior to discharge. Because of the high incidence of neural hearing loss associated with significantly elevated bilirubin, these infants should be referred for audiological assessment to include ABR measures.

## **Follow-up and Case Management**

EHDI is part of a continuum of care that progresses from screening to assessment to amplification (if elected) to educational intervention. Assessment referrals need to be made to audiologists with expertise in pediatric physiological and behavioral assessment and management. See the Minnesota Department of Health (MDH) website for Audiology Regional Centers that provide pediatric diagnostic assessments and habilitation services in Minnesota ([www.health.state.mn.us/newbornscreening](http://www.health.state.mn.us/newbornscreening)).

The National recommended time line for hearing screening and follow-up includes: complete screening no later than one month of age, assessment no later than three months of age, and amplification (if elected) and intervention no later than six months of age. With prompt referral and follow-up, Minnesota children will receive appropriate care and services even earlier.

### **Middle Ear Effusion**

Although persistent middle ear effusion necessitates medical referral, which might delay the evaluation timeline several weeks, diagnostic audiological evaluation must not be repeatedly postponed solely due to middle ear dysfunction and should be completed before 3 months of age. The information from the diagnostic audiological evaluation is valuable both in determining the extent of the effect of the middle ear condition on the infant's hearing, and identifying whether an underlying sensorineural

hearing loss exists, thereby impacting the course of both medical and educational intervention.

**Positive Risk Factor** (JCIH 2007 clarification document)

The timing and number of hearing re-evaluations for children with risk factors (Appendix A) should be customized and individualized depending on the relative likelihood of a subsequent delayed-onset hearing loss. Infants who pass the neonatal screening but have a risk factor should have at least 1 diagnostic audiology assessment by 24 to 30 months of age. Early and more frequent assessment may be indicated for children with cytomegalovirus (CMV) infection, syndromes associated with progressive hearing loss, neurodegenerative disorders, trauma, or culture-positive postnatal infections associated with sensorineural hearing loss; for children who have received ECMO or chemotherapy; and when there is caregiver concern or a family history of hearing loss.

Without an adequate follow-up plan, even the best EHDI program is ineffective. Refer to the current Minnesota Department of Health (MDH) Infant [Assessment](#) and Pediatric [Amplification](#) Guidelines.

## QUALITY ASSURANCE

Components of a quality assurance program include data management, screener performance, site performance, outcome measures, and follow-up compliance. The goal of quality assurance is information management and accountability to the following stakeholders:

- Family and the infant
- Screeners
- EHDI manager
- Clinical and educational audiologist
- Primary Care Provider (Medical Home)
- Hospital
- State of Minnesota
- Otolaryngologists
- Parents
- Medical and Educational Specialists
- Advocates

## Appendix A

### **RISK INDICATORS ASSOCIATED WITH PERMANENT CONGENITAL, DELAYED-ONSET, OR PROGRESSIVE HEARING LOSS IN CHILDHOOD**

Risk indicators that are marked with a "§" are of greater concern for delayed-onset hearing loss.

1. Caregiver concern§ regarding hearing, speech, language, or developmental delay.
2. Family history§ of permanent childhood hearing loss.
3. All infants with or without risk factors requiring neonatal intensive care for greater than 5 days, including any of the following: ECMO,§ assisted ventilation, exposure to ototoxic medications (gentamicin and tobramycin) or loop diuretics (furosemide/Lasix). In addition, regardless of length of stay: hyperbilirubinemia requiring exchange transfusion.
4. In utero infections, such as CMV,§ herpes, rubella, syphilis, and toxoplasmosis.
5. Craniofacial anomalies, including those that involve the pinna, ear canal, ear tags, ear pits, and temporal bone anomalies.
6. Physical findings, such as white forelock, that are associated with a syndrome known to include a sensorineural or permanent conductive hearing loss.
7. Syndromes associated with hearing loss or progressive or late-onset hearing loss,§ such as neurofibromatosis, osteopetrosis, and Usher syndrome; other frequently identified syndromes include Waardenburg, Alport, Pendred, and Jervell and Lange-Nielson.
8. Neurodegenerative disorders,§ such as Hunter syndrome, or sensory motor neuropathies, such as Friedreich ataxia and Charcot-Marie-Tooth syndrome.
9. Culture-positive postnatal infections associated with sensorineural hearing loss,§ including confirmed bacterial and viral (especially herpes viruses and varicella) meningitis.
10. Head trauma, especially basal skull/temporal bone fracture§ that requires hospitalization.
11. Chemotherapy.§

\* *Risk factor references available in JCIH 2007 Position Statement*

## Appendix B

### Hearing & Speech Milestones

Birth to 3 months	<ul style="list-style-type: none"><li>• startles to loud sounds</li><li>• quiets or smiles when spoken to</li><li>• seems to recognize your voice and quiets if crying</li><li>• increases or decreases sucking behavior in response to sound</li><li>• makes pleasure sounds (cooing, gooing)</li><li>• cries differently for different needs</li><li>• smiles when sees you</li></ul>
3 to 6 months	<ul style="list-style-type: none"><li>• looks or turns toward a new sound</li><li>• responds to changes in tone of voice</li><li>• vocalizes excitement and displeasure</li><li>• enjoys rattles and other toys that make sounds</li><li>• babbling sounds more speech-like (such as ooh, aah, and ba-ba)</li></ul>
6 to 12 months	<ul style="list-style-type: none"><li>• enjoys games like peek-a-boo and pat-a-cake</li><li>• responds to his/her own name, telephone ringing, someone's voice, even when not loud</li><li>• knows words for common things (cup, shoe) and sayings ("bye-bye")</li><li>• imitates different speech sounds</li><li>• starts to respond to requests such as "come here"</li><li>• looks at things or pictures when someone talks about them</li><li>• has one or two words (bye-bye, dada, mama). although they may not be clear</li></ul>

## Appendix C

### HEARING SCREENING RESULT AND FOLLOW-UP PROCESS

(a) Pass Both Ears, No Risk Indicators

- ✓ Passes screen in both ears (AABR or OAE)
- ✓ Does not have any risk indicators for hearing loss (see risk indicators in Appendix A.)
- ✓ Parents receive information (oral and written, culturally appropriate) regarding developmental milestones for auditory, speech, and language skills. In addition, parents should be encouraged to return for audiologic evaluation if their child is not meeting these milestones, or if they have any concerns about their child's hearing at any point in the future.

(b) Pass Both Ears, Positive Risk

- ✓ Passes screen in both ears (AABR or OAE)
- ✓ Infant has at least one risk indicator for hearing loss (see risk indicators for hearing loss, Appendix A, at the end of this document).
- ✓ Infant is referred and scheduled for repeat hearing screen at three or six months of age with audiologist.
- ✓ If infant's hearing screen is normal at age three months, a complete audiological evaluation, using visual reinforcement audiometry (VRA), should be scheduled for six months developmental age.
- ✓ If follow-up screening is abnormal at three or six months, diagnostic ABR is performed as soon as feasible.
- ✓ Parents receive information (oral and written, and culturally appropriate) regarding developmental milestones for auditory, speech, and language skills. In addition, parents should be encouraged to return for audiologic evaluation if their child is not meeting these milestones, or if they have any concerns about their child's hearing at any point in the future.

(c) Does Not Pass in One or Both Ears, or Incomplete Test

- ✓ Does not pass AABR or OAE hearing screen in one or both ears.
- ✓ Testing cannot be completed in one or both ears prior to discharge or as an outpatient re-screen.
- ✓ If repeat hearing screening is abnormal for one or both ears, the infant is referred and scheduled for diagnostic audiological evaluation as soon as feasible. National standards specify diagnostic testing needs to occur no later than three months of age.

## References

American Speech-Language-Hearing Association (2004). *Guidelines for the Audiologic assessment of children from birth to 5 years of age*. Rockville, MD: ASHA.

CDC EHDI National Goals and Objectives, Final Version by the EHDI Data Committee, July 13, 2006.

Gorga, M.P., Neely, T.S., Ohlrich, B., Hoover, B., Redner, J. & Peters, J. (1999). From laboratory to clinic: A large scale study of distortion product otoacoustic emissions in ears with normal hearing and ears with hearing. *Ear and Hearing*, 18, 440-455.

Hall, J.W., Smith, S.D., & Popelka, G.R. (2004). Newborn hearing screening with combined otoacoustic emissions and auditory brainstem responses. *Journal of the American Academy of Audiology*, 15, 414-425.

HRSA, Final Report. Evaluation of Universal Newborn Hearing Screening and Intervention Program. Based on 2005 – 2006 Data.

Joint Committee on Infant Hearing Position Statement (2007), *Year 2007 position statement: Principles and guidelines for early hearing detection and intervention*. *Pediatrics*, 120, 898-921.

Joint Committee on Infant Hearing Position Statement (2007) Update. *Clarification for Year 2007 JCIH Position Statement*. Retrieved May 2008, from [www.jcih.org/Clarification%20Year%202007%20statement.pdf](http://www.jcih.org/Clarification%20Year%202007%20statement.pdf)

Minnesota Statute 144.966. Early Hearing Detection and Intervention Program. 2007. Retrieved from [www.leg.state.mn.us/leg/statutes](http://www.leg.state.mn.us/leg/statutes).

National Consortium for Newborn Hearing Screening. (Nov 16-18, 1995). *TEOAE-based universal newborn hearing screening*. Georgetown University School of Medicine, Washington DC.

Norton, S., Gorga, M., Widen, J., Folsom, R., Sininger, Y., Cone-Wesson, B., Vohr, B., & Fletcher, K. (2000). Identification of neonatal hearing impairment: A multicenter investigation. *Ear and Hearing*, 21 (5), 348-356.

Ontario Health and Long-Term Care Ministry—Infant Hearing Program (2002). *Universal infant hearing screening assessment and communication development: Local implementation support document*.

Washington State Department of Health Protocol for Newborn Hearing Screening. Retrieved June 24, 2007, from [http://www.doh.wa.gov/cfh/mch/Genetics/ehddi/Documents/newborn\\_protocol.pdf](http://www.doh.wa.gov/cfh/mch/Genetics/ehddi/Documents/newborn_protocol.pdf)

*Revised from the MDH Audiology Task Force, Protocols for Organization and Administration of UNHS Programs, Document (2005.)*