

Adolescent Mental Health and/or Depression Screening Measure Specifications 2023 Measurement Year

Summary of Changes	NONE		
Description	The percentage of patients 12-17 years of age who were screened for mental health and/or depression		
Measurement Period	FINAL 2023 MY: January 1, 2023 through December 31, 2023		
	PRELIMINARY 2024 MY: January 1, 2024 through December 31, 2024		
Eligible Population	Eligible Specialties	Family Medicine, Internal Medicine, Pediatric/Adolescent Medicine	
	Eligible Providers	Medical Doctor (MD), Doctor of Osteopathy (DO), Physician Assistant (PA), Advanced Practice Registered Nurses (APRN)	
	Ages	12 years or older at the start of the measurement period AND less than 18 years at the end of the measurement period	
	Event	At least one office well child visit or eligible telehealth encounter (<i>Adolescent Well Child Visit</i> Value Set) performed or supervised by an eligible provider in an eligible specialty during the measurement period	
Denominator	The eligible population		
Numerator	The number of patients in the denominator who completed ANY of the following mental health and/or depression screening tools during the measurement period AND have the screening tool result documented in the medical record:		
	• Patient Health Questionnaire – 9 item version (PHQ-9)		
	PHQ-9M Modified for Teens and Adolescents		
	Kutcher Depression Scale (KADS)		
	Beck Depression Inventory II (BDI-II)		
	Beck Depression Inventory Fast Screen (BDI-FS)		
	Child Depression	Inventory (CDI)	
	Child Depression Inventory II (CDI-2)		
	• Patient Health Questionnaire – 2 item version (PHQ-2)		
	• Pediatric Symptom Checklist – 17 item version (PSC-17) - parent version		
	• Pediatric Symptom Checklist – 35 item (PSC-35) - parent version		
	• Pediatric Symptom Checklist – 35 item Youth Self-Report (PSC Y-SR)		
	Global Appraisal of Individual Needs screens for mental health and substance abuse (GAIN-SS)		



Adolescent Mental Health and/or Depression Screening Measure Specifications 2023 Measurement Year

Required Exclusions	The following exclusions must be applied to the eligible population:		
	• Patient had a diagnosis of schizophrenia (<i>Schizophrenia or Psychotic Disorders</i> Value Set) any time prior to the end of the measurement period		
	• Patient had a diagnosis of bipolar disorder (<i>Bipolar Disorder</i> Value Set) any time prior to the end of the measurement period		
	• Patient had an active diagnosis of depression or dysthymia (<i>Major Depression or Dysthymia</i> Value Set) any time during the measurement period		
	• Patient had a diagnosis of personality disorder (<i>Personality Disorder</i> – <i>Emotionally Labile</i> Value Set) any time prior to the end of the measurement period		
	• Patient had moderate, severe, or profound intellectual disabilities (<i>Intellectual Disability</i> Value Set) any time prior to the end of the measurement period		
Measure Scoring	Rate/Proportion		
Interpretation of Score	Higher score indicates better quality		
Measure Type	Process		