

# Meningococcal Vaccination Pocket Guide

## Meningococcal Vaccines

Meningococcal serogroup(s) involved	ACWY	B
Recommended age indications*	<ul style="list-style-type: none"> <li>▪ At risk age 2 months and older</li> <li>▪ Routine adolescent:                             <ul style="list-style-type: none"> <li>▪ 11-12 years</li> <li>▪ Booster at 16 years</li> </ul> </li> <li>▪ At risk adults</li> </ul>	<ul style="list-style-type: none"> <li>▪ At risk 10 years and older</li> <li>▪ Consider for anyone 16-23 years</li> </ul>
Vaccine names and associated brand	MenACWY-crm:    Menveo MenACWY-D:     Menactra	MenB-4C:        Bexsero MenB-FHbp:     Trumenba
Immunization requirement?**)	Yes, grades 7-12	No

\*Recommended by the Advisory Committee on Immunization Practices (ACIP), see FDA-approved package insert for licensed age indication and dosage schedule.

\*\*Documentation required showing receipt of meningococcal vaccine or a medical or non-medical exemption.

## Risk Indications for Meningococcal Vaccines

Risk indications	Meningococcal ACWY (MenACWY, MCV4)	Meningococcal B (MenB)
Complement component deficiency	Yes	Yes
Anatomic / functional asplenia	Yes	Yes
HIV infected	Yes	No
Travel	Yes	No
Microbiologist working with <i>N. Meningitidis</i>	Yes	Yes
First-year college student living in residence hall	Yes	No
Military recruit	Yes	No
Community outbreak	Yes, for outbreaks involving serogroups A, C, W, Y	Yes, for outbreaks involving serogroup B

### Key things to remember

- When giving the first dose of MenACWY, remind parent and pre-teen that a booster is needed at age 16.
- MenACWY and MenB can be given at the same visit.
- Use the same MenB product for all doses of the series.

### More information

Find additional resources and guidance on the Meningococcal Disease Information for Health Care Professionals ([www.health.state.mn.us/divs/idepc/diseases/meningococcal/hcp.html](http://www.health.state.mn.us/divs/idepc/diseases/meningococcal/hcp.html)) website.