

Table 2c. Number of Cases and Rates (per 100,000 persons) of Primary/Secondary Syphilis by Residence, Age, and Gender¹-- Minnesota, 2016

Group	Males		Females		Total ^v		
	Cases	%	Cases	%	Cases	%	Rate
Residence							
Minneapolis	111	42%	14	38%	127	42%	33.2
St. Paul	27	10%	2	5%	29	9%	10.2
Suburban ¹	84	31%	10	27%	94	31%	4.3
Greater Minnesota	45	17%	11	30%	56	18%	2.3
Age							
< 15 yrs	0	0%	1	3%	1	0%	0.1
15-19 yrs	9	3%	5	14%	14	5%	3.8
20-24 yrs	44	16%	5	14%	49	16%	13.8
25-29 yrs	42	16%	6	16%	49	16%	13.1
30-34 yrs	45	17%	14	38%	60	20%	17.5
35-39 yrs	37	14%	1	3%	38	12%	11.6
40-44 yrs	33	12%	0	0%	33	11%	9.4
45-49 yrs	17	6%	3	8%	20	7%	4.9
50-54 yrs	23	9%	1	0%	24	8%	6.0
55+ yrs	17	6%	1	3%	18	6%	1.4
Race/Ethnicity							
White, Non-Hispanic	145	54%	15	41%	162	53%	3.5
Black, Non-Hispanic	59	22%	12	32%	71	23%	25.3
American Indian	10	4%	7	19%	17	6%	25.3
Asian/PI	18	7%	1	3%	19	6%	8.6
Other ^{ii, iii}	0	0%	0	0%	0	0%	x
Unknown ⁱⁱⁱ	10	4%	0	0%	10	3%	x
Hispanic ^{iv}	25	9%	2	5%	27	9%	10.8
Total	267		37		306		5.8

Notes

Data exclude cases diagnosed in federal or private correctional facilities.

U.S. Census 2010 data is used to calculate rates.

^I Suburban is defined as the seven-county metropolitan area (Anoka, Carver, Dakota, Hennepin, Ramsey, Scott and Washington Counties, excluding the cities of Minneapolis and St. Paul).

^{II} Includes persons reported with more than one race.

^{III} No comparable population data available to calculate rates.

^{IV} Persons of Hispanic origin may be of any race.

^V Total includes 2 cases of primary/secondary syphilis diagnosed in transgendered persons (male to female).

Minnesota Department of Health
STD, HIV and TB Section
(phone) 651-201-5414
www.health.state.mn.us

04/19/2017



To obtain this information in a different format, call: 651-201-5414.