

Dawaynta Loogu Taloglaay Caabuqa Qaaxada (TB) Qarsoon: Moxifloxacin

Baaritaankaagu waxa uu muujinayaa inaad qabto caabuqa TB-da qarsoon, sidoo kale loo yaqaan "LTBI". Caabuqa TB-da Qarsoon macnaheedu waa jeermiska TB ayaa ku jira jirkaaga laakiin waxay u egtay jeermis hurda. Jeermiska TB-da qarsoon kuma waxyeelaynayo adiga oo umana gudbi karo dadka kale. Haddii jeermiska TB-du uu tooso oo uu noqdo mid firfircoon, markaa waxay kuu keeni kartaa jirro. Tani waxa lagu magacaabaa cudurka TB-da firfircoon. Waxad u gudbin kartaa TB-da dadka kale haddii aad qabto cudurka TB-da firfircoon. Qaadashada dawooyinka loogu talagalay caabuqaga TB-da qarsoon ayaa kaa caawin karta inaad kahortagto jeermiska TB-da qarsoon inuu kugu soo kaco oo uu noqdo cudurka TB-da firfircoon.

Jeermiska qaaxada kugu dhacday ayaa ahayd qaaxada dawoyinka badan iska caabbisa (MDR TB). Qaaxada MDR waa marka jeermiska TB-du uu iska caabbiyo ugu yaraan laba ka mida kuwa ugu waaweyn dawooyinka qaaxada. Jeermiska xooggan laguma dawayn karo dawooyinka aan caadi ahaan u isticmaalno inaan ku dawaynno LTBI. Sababtan darteed waxaad u baahan doontaa dawo ka duwan ta qofka qaba LTBI ee caadigaah.

Sideen uga hortagi karaa cudurka Qaaxada firfircoon?

Waxad qaadan kartaa dawo si aad uga hortagto inuu kugu dhaco cudurka qaaxada firfircoon. Moxifloxacin waa dawo loo isticmaalo in lagu daweeyo LTBI. Waxay dishaa jeermiska qaaxada ee hurda kahor intaanay jirro kugu keenin. Waxay qaadan kartaa bilo badan dawada inay ku disho jeermiska qaaxada maxaayeelay jeermiska ayaa xooggan. Qaado Moxifloxacin inta goor iyo illaa inta dakhtarkaaga ama kalkaalisadaadu ay ku farto.

Maxaan dawada u qaataa haddii aanan dareemayn jirro?

Jeermiska TB-da hurda ayaa aad u fudud in la dilo kahor intaanay toosin ee aanay kugu keenin jirro.

Maxaa dhacaya haddii aanaan qaadan daawada?

Haddii aanad qaadan Moxifloxacin, aad ka tagto maalmo aad u badan, ama aad joojiso qaadashada dawada kahor dakhtarkaaga ama kalkaalisadaadu intaanay ku dhihin sidaas, waxa laga yaabaa inaad la jirrato Qaaxada firfircoon. Waa muhiim in la dilo jeermiska si markaa adiga iyo qoyskaagu uu caafimaad qabo.

Ka warran haddii aan qaatay tallaalka BCG?

BCG waxay ka ilaalisaa carruurta qaaxada laakiin kaliya dhawr sanno. Kama ilaalin karto qaadista LTBI ama cudurka qaaxada firfircoon. Haddii aad qaadatay BCG, waa inaad wali qaadataa dawada loogu talagalay LTBI si aad u ilaaliso naftaada iyo dadka kalaba.

Ka warran haddii aanaan iska bixin karin lacagta Moxifloxacin?

Waydii dhakhtarkaaga ama kalkaalisada si ay kaaga caawiso inaad ka hesho dawada Qaaxada Barnaamijka Qaaxada ee Waaxda Caafimaadka Minnesota.



Ka warran haddii aan guuro?

U sheeg dhakhtarkaaga ama kalkaalisada kahor intaadan u guurin gobol ama magaalo kale. Waxay kaa caawin karaan inaad sii wadato daawada qaaxada kadib markaad guurto.

Maxay tahay inaan ka oggaado arrinta ku saabsan daawada LTBI?

- U sheeg dhakhtarkaaga wixii ku saabsan daawooyinka kale ee aad qaadato.
- Ha qaad Moxifloxacin 2 saacadood gudahood marka aad cunto ama aad cabto wax kasta oo caanoleh ama samaysan, dawooyinka laab-jeexa (antacids) ama maltifiitaamiinnada (multivitamins).
- Haddii aad sonkorow qabtid, fiiri heerka sonkorta dhiiggaaga si joogto ah.
- Iska ilaali inaad uur yeelato inta aad qaadaneys Moxifloxacin. Wac dhakhtarkaaga ama kalkaalisada islamarkiiba haddii aad uur leedahay.

Dadka intooda ugu badan waxay qaataan Moxifloxacin dhibaato la'aan. Dadka qaar waxay yeeshaan dhibaatooyin, sida:

- Calool majiir ama matag
- Madax xanuun ama dawakhaad (madaxa wareer)
- Shuban daran
- Calool xanuun
- Finan maqaarka ah ama cuncun
- Maqaarka iyo indhaha oo huruud ah
- Xanuun, gubasho ama xoqid gacmaha ama lugaha
- Xanuunka muruqyadaada ama xubnahaaga
- Dareemi niyada-xumo ama isbeddelka niyadda

Haddii aad yeelato mid ka mid ah dhibaatooyinkan:

1. Jooji qaadashada daawadaada.
2. Islamarkiiba wac dhakhtarkaaga ama kalkaalisada. Ha sugin ballantaada xigta.

Minnesota Department of Health
Tuberculosis (TB) Prevention and Control Program
P.O. Box 64975, St. Paul, MN 55164-0975
651-201-5414 | www.health.state.mn.us/tb

Si aad ugu hesho macluumaadkan qaab kale, wac:
651-201-5414. Waxaa lagu daabacay warqad dib loo rogay.



Xusuusnow inaad qaadato daawadaada qaaxada.

Haddii aad seegto maalmo badan, daawadu ma shaqeyn karto. Halkan waxaa ku yaal dhowr siyaabood oo aad ku xasuusan kartid:

- Dhig Moxifloxacin meel aad ka arki doonto.
 - Qaado dawadaada Moxifloxacin isla waqti maalinta. Tusaale ahaan, ka dib markaad cadayato ilkahaaga ama ka hor intaanad seexan.
 - Weydiiso qof inuu ku xasuusiyo.
 - Ku calaamade jadwalkaaga ka dib markaad qaadatid Moxifloxacin.
 - Ku samayso xasuusin ama digniin taleefankaaga gacanta.
 - Isticmaal sanduuqa xusuusinta kiniinnada.
- Haddii aad seegto maalmo, u qor si aad ugu sheegto dhakhtarkaaga ama kalkaalisadaada ballanta soo socota.

Oofidhammaan ballamahaaga caafimaadka.

Dhakhtarkaaga ama kalkaalisada ayaa kaa caawin doona inaad hubiso inay dawayntaadu si fiican u socoto.

Ballantayda soo socota ee daawada qaaxada (TB)

Taariikhda:

Wakhtiga:

Dakhtarka ama kaaalisada:

Goobta: